# ACSM CEP Clinical Exer Prep

Free Practice Test — 30 Real Exam-Style Questions

with full answer key & explanations

**Unlock the full bank of 710 questions**
**+ unlimited timed mock exams + mistake book**

**Practice on the web:** https://certs.theorypractice.app/cepp
$2.99 / week · $6.99 / month · cancel anytime

**What you unlock: all 710 questions • unlimited timed mock exams • mistake book • instant explanations**

**Study offline on the free app — search your exam on the App Store or Google Play**

## Practice Questions

Try all 30 first, then check the answer key at the back.

**Want the other 680+ questions & full timed mock exams? Unlock at https://certs.theorypractice.app/cepp**

**1. Which training scenario is most likely to cause Delayed Onset Muscle Soreness (DOMS)?**

A. Low-intensity cycling
B. Moderate-intensity swimming
C. Yoga stretching session
D. High-intensity downhill running

**2. Which of the following individuals is most likely to experience symptoms of exercise burnout?**

A. An amateur cyclist who takes rest days and varies his training routine.
B. A yoga enthusiast who practices yoga daily but includes different yoga styles.
C. A professional triathlete who trains for multiple hours every day without adequate rest days.
D. A recreational runner who jogs 3 times a week for 30 minutes.

**3. For a tennis player, which of the following would not be a necessary component of their exercise prescription to address the demands of their sport?**

A. Shoulder strengthening exercises
B. Endurance training
C. High-intensity interval training
D. Agility drills

**Study offline on the free app — search your exam on the App Store or Google Play**

**4. A patient reports that while running the other day, he felt a "snap" in his left calf. Ever since, he's noticed severe pain and difficulty bearing weight on that leg. What happened to this patient and how should the Clinical Exercise Physiologist (CEP) respond?**

A. This patient is possibly overreacting to the experience of feeling a "snap" and should continue his exercise regimen without any modifications.
B. This patient may have a minor calf strain that should resolve with rest. He can resume light exercise after a day or two.
C. This patient might have an issue with his calf muscle, but he should continue to exercise and just avoid movements that cause pain.
D. This patient probably suffered an Achilles tendon rupture. He should be encouraged to follow up with his doctor before continuing exercise.

**5. A client tells her CEP that she used to enjoy playing tennis and wants to get back into the game. What exercises would best help the client achieve this specific goal of returning to playing tennis?**

A. Cycling and running long distances
B. Heavy weightlifting with minimal repetitions
C. Rotational core exercises, leg agility drills, shoulder mobility exercises
D. Swimming laps

**6. Which of the following is true of patients with chronic obstructive pulmonary disease (COPD) regarding exercise in high-altitude environments?**

A. High-altitude air has no effect on exercise tolerance for most patients with COPD
B. Exercise in high-altitude environments could more easily trigger a COPD exacerbation
C. Exercise in high-altitude environments tends to be more comfortable for most patients with COPD
D. Patients with COPD should never exercise in high-altitude environments

**Want the other 680+ questions & full timed mock exams? Unlock at https://certs.theorypractice.app/cepp**

**7. Which of the following is a crucial factor to consider when prescribing exercise for a patient post-cardiac bypass surgery (CBS)?**

A. Patients who have undergone CBS typically do not require any exercise modifications
B. Patients may need up to a year or longer to achieve full recovery from cardiac bypass surgery
C. High-intensity exercises are important to incorporate early in the rehabilitation process
D. Patients who have undergone CBS tend to have worse outcomes than those who have had other types of heart surgery

**Unlock all 710 questions + timed mock exams**
→ https://certs.theorypractice.app/cepp
$2.99/week or $6.99/month · cancel anytime · scan to start
Unofficial study material · not affiliated with any certifying body

**8. Which of the following is not a component of the cardiovascular fitness assessment?**

A. Blood pressure measurement
B. Agility test
C. VO2 max test
D. Heart rate monitoring

**9. Considering the long-term health benefits of strength and resistance training in seniors, which of the following statements is correct?**

A. Strength and resistance training cannot improve muscle mass, only bone density can be improved in seniors.
B. Seniors should avoid strength and resistance training due to high risk of injury and minimal health benefits.
C. Strength and resistance training can significantly improve muscle mass and bone density in seniors even if started late in life.
D. Strength and resistance training must be started in youth to have any significant health benefits in old age.

**Study offline on the free app — search your exam on the App Store or Google Play**

**10. Which of the following regarding exercise and hypertension is true?**

A. Only high-intensity interval training (HIIT) is beneficial for reducing blood pressure.
B. Regular aerobic exercise can significantly reduce blood pressure in hypertensive patients.
C. Strength training increases blood pressure and is not recommended for hypertensive patients.
D. Exercise has no significant impact on blood pressure for hypertensive patients.

**11. Which of the following is a characteristic of a patient showing minimal improvement in blood pressure despite a consistent hypertensive exercise program?**

A. A patient whose overall cardiovascular fitness improves, but reports that they do not enjoy the exercise sessions
B. A patient whose blood pressure remains essentially unchanged after six months of structured exercise targeted at hypertension management
C. A patient whose systolic blood pressure decreases by 15 mmHg after two months of hypertensive exercise program
D. A patient whose blood pressure initially drops but then rises back to pre-exercise levels after discontinuing the program for two weeks

**Unlock all 710 questions + timed mock exams**
→ **https://certs.theorypractice.app/cepp**
$2.99/week or $6.99/month · cancel anytime · scan to start
Unofficial study material · not affiliated with any certifying body

**12. Which of the following is true of Exercise-induced Asthma?**

A. Exercise-induced Asthma is primarily a disease of the digestive system.
B. Exercise-induced Asthma is twice more common in men than in women.
C. Patients with Exercise-induced Asthma can manage their symptoms through proper warm-up routines and medication.
D. Exercise-induced Asthma symptoms are typically triggered by exposure to cold air alone.

**Want the other 680+ questions & full timed mock exams? Unlock at https://certs.theorypractice.app/cepp**

**13. Which physiological improvements are most commonly observed in patients undergoing regular aerobic exercise?**

A. Reduced bone density, increased risk of joint injuries, and increased fatigue
B. Increased muscle atrophy, worsened flexibility, and decreased metabolic rate
C. Increased blood pressure, decreased cardiac output, and reduced mental health
D. Improved cardiovascular function, increased lung capacity, enhanced overall endurance

**14. During the eccentric phase of a bicep curl, what is the motion at the elbow?**

A. Extension
B. Flexion
C. Abduction
D. Pronation

**15. Which of the following would be a suitable balance exercise consideration for an older adult at risk of falls?**

A. Advise the client to run on a treadmill
B. Encourage the client to lift heavy weights overhead
C. Use loud music to cue balance adjustments
D. Have the client practice standing on one foot while holding onto a chair for support

**Study offline on the free app — search your exam on the App Store or Google Play**

**16. In the context of a clinical exercise setting, which of the following plays a crucial role in providing support and cushioning for joints?**

A. Tendons
B. Cartilage
C. Ligaments
D. Muscles

**Unlock all 710 questions + timed mock exams**
→ https://certs.theorypractice.app/cepp
$2.99/week or $6.99/month · cancel anytime · scan to start
Unofficial study material · not affiliated with any certifying body

**17. Which of the following is true of plantar fasciitis?**

A. It is commonly referred to as "runner's knee"
B. It is an inflammation of the Achilles tendon
C. It is an acute injury that can occur from a single event
D. It is commonly referred to as "heel spur syndrome"

**18. In the context of weightlifting, which of the following best describes a static hold exercise?**

A. An exercise where a particular muscle group is held in a fixed position, producing force without causing movement at the joint
B. An exercise characterized by rapid and explosive movements like cleans or snatches
C. An exercise involving movements at a constant speed, often requiring specialized machines
D. An exercise that alternates between lifting and lowering a weight with controlled, steady motion

**Want the other 680+ questions & full timed mock exams? Unlock at https://certs.theorypractice.app/cepp**

**19. A patient has a resting heart rate of 65 beats per minute (bpm). How should the CEP interpret this finding?**

A. This is a normal resting heart rate.
B. This heart rate falls within the "dangerously low" range. The patient should be referred for extensive medical testing before starting exercise.
C. This heart rate is indicative of tachycardia.
D. This heart rate may cause a life-threatening situation for the client. The CEP should contact emergency services immediately.

**20. Which of the following symptoms in a client would necessitate an immediate referral to their healthcare provider?**

A. Occasional dizziness when standing up quickly
B. Minor lower back pain after lifting weights
C. Sudden onset of chest pain, shortness of breath, and lightheadedness
D. Mild soreness in the knees after a jogging session

**21. During a high-intensity workout, what is the term used to describe the state when muscle cells lack sufficient oxygen?**

A. Hypoxia
B. Hypercapnia
C. Anemia
D. Ischemia

**Unlock all 710 questions + timed mock exams**
→ https://certs.theorypractice.app/cepp
$2.99/week or $6.99/month · cancel anytime · scan to start
Unofficial study material · not affiliated with any certifying body

**Study offline on the free app — search your exam on the App Store or Google Play**

**22. When instructing a patient on how to use a peak flow meter, which of the following is the most accurate method?**

A. Ask the patient to take a deep breath, place the mouthpiece into the mouth, and blow out as hard and fast as possible into the device.
B. Ask the patient to exhale slowly and steadily into the device after a normal breath.
C. Ask the patient to inhale as fast as possible through the device.
D. Ask the patient to take a deep breath, place the mouthpiece into the mouth, and blow out steadily and slowly into the device.

**23. According to the Health Belief Model, which factors influence an individual's likelihood to engage in preventive health behaviors?**

A. Perceived barriers only
B. Cues to action only
C. Self-efficacy only
D. Perceived susceptibility, perceived severity, perceived benefits

**24. Which of the following behaviors would indicate that a patient is in the contemplation stage of adopting a healthier diet?**

A. The patient has maintained a healthy diet for over six months.
B. The patient has begun to include one or two healthy snacks each day.
C. The patient states that they are not interested in changing their diet.
D. The patient has started asking about different types of healthy meal plans.

**Want the other 680+ questions & full timed mock exams? Unlock at https://certs.theorypractice.app/cepp**

**25. Which of the following is a characteristic of exercise addiction?**

A. Gaining significant weight due to inactivity
B. Only exercising in social settings
C. Continuing to exercise despite injury
D. Showing no interest in physical activity

**Unlock all 710 questions + timed mock exams**
→ https://certs.theorypractice.app/cepp
$2.99/week or $6.99/month · cancel anytime · scan to start
Unofficial study material · not affiliated with any certifying body

**26. For a patient with diabetic peripheral neuropathy (DPN), what is a critical safety consideration during exercise testing?**

A. This patient will never recover the sensation in their feet.
B. This patient may have an increased risk of falls due to decreased foot sensation.
C. This patient will likely exhibit a significant reduction in upper body muscular strength.
D. This patient will probably encounter severe difficulties in cognitive functions during exercise testing.

**27. How does the Naughton Protocol differ from the modified Bruce Protocol?**

A. The Naughton Protocol is designed for a more gradual increase in intensity, with lower starting speeds and fewer changes in speed and grade compared to the modified Bruce Protocol.
B. The Naughton Protocol starts at a higher speed compared to the modified Bruce Protocol.
C. The modified Bruce Protocol decreases speed and grade gradually, while the Naughton Protocol increases them rapidly.
D. The Naughton Protocol includes rest periods between stages, unlike the modified Bruce Protocol.

**Study offline on the free app — search your exam on the App Store or Google Play**

**28. During a graded exercise test, which physiological response best indicates the achievement of maximal aerobic capacity?**

A. An increase in blood pressure without a plateau
B. A decrease in respiratory rate as workload increases
C. A plateau in oxygen consumption ($$\dot{V}O_2$$) despite an increase in workload
D. A steady increase in heart rate with increasing workload

**29. When conducting a graded exercise test (GXT), which of the following could lead to an inaccurate assessment of cardiovascular fitness?**

A. Incorrect calibration of the treadmill or ergometer
B. Gradual increase of workload according to the established protocol
C. Monitoring heart rate using a heart rate monitor
D. Ensuring the participant is familiar with the exercise equipment

**30. Which of the following statements is false regarding the physiology during a maximal exercise test?**

A. Lactic acid levels rise significantly after the anaerobic threshold is reached.
B. The VO2 max is achieved when oxygen consumption plateaus despite increased workload.
C. The respiratory exchange ratio (RER) typically exceeds 1.0 at maximal exertion.
D. The heart rate remains stable after reaching the anaerobic threshold.

**Unlock all 710 questions + timed mock exams**
→ **https://certs.theorypractice.app/cepp**
$2.99/week or $6.99/month · cancel anytime · scan to start
Unofficial study material · not affiliated with any certifying body

## Answer Key & Explanations

**You just practised 30 of 710. Unlock every question + timed mocks at https://certs.theorypractice.app/cepp**

**1. D — High-intensity downhill running**
Answer: High-intensity downhill running Eccentric contractions, such as those occurring during high-intensity downhill running, significantly contribute to muscle damage and Delayed Onset Muscle Soreness (DOMS). This kind of activity involves the lengthening of muscle fibers under tension, leading to microtears and subsequent soreness. While other exercises like yoga or low-intensity activities can cause muscle soreness, they are less likely to cause DOMS compared to high-intensity eccentric exercises.

**2. C — A professional triathlete who trains for multiple hours every day without adequate rest days.**
Answer: A professional triathlete who trains for multiple hours every day without adequate rest days. Exercise burnout is a result of consistently high intensity and volume of training without sufficient recovery time. Athletes who balance intense training sessions with proper rest and varied routines are less likely to experience burnout. Recreational runners, amateur cyclists who take rest days, and yoga enthusiasts who practice different styles are less likely to face burnout because they allow for recovery and variation in their activities.

**3. C — High-intensity interval training**
Answer: High-intensity interval training While high-intensity interval training (HIIT) might be beneficial in general fitness programs, it is not critical for a tennis player's specific needs. Agility drills are essential for quick directional changes; shoulder strengthening exercises are important due to the repetitive overhead motion; and endurance training ensures stamina throughout matches.

**4. D — This patient probably suffered an Achilles tendon rupture. He should be encouraged to follow up with his doctor before continuing exercise.**
Answer: This patient probably suffered an Achilles tendon rupture. He should be encouraged to follow up with his doctor before continuing exercise. This patient is likely experiencing symptoms of an Achilles tendon rupture, which often presents as a sudden snap followed by severe pain and difficulty bearing weight. It is crucial to have this injury evaluated by a physician to confirm the diagnosis and discuss treatment options, which may include surgery. The patient should avoid any weight-bearing activity until medically cleared.

**5. C — Rotational core exercises, leg agility drills, shoulder mobility exercises**
Answer: Rotational core exercises, leg agility drills, shoulder mobility exercises. Playing tennis requires good rotational core strength for swings, leg agility for quick movements, and shoulder mobility to prevent injuries. Although other forms of exercise can improve fitness, they do not specifically target the skills and strengths needed for tennis.

**6. B — Exercise in high-altitude environments could more easily trigger a COPD exacerbation**
Answer: Exercise in high-altitude environments could more easily trigger a COPD exacerbation Patients with COPD should always ensure that they have their rescue inhalers available when exercising in high-altitude environments. The low oxygen levels can trigger a bronchospasm and exacerbate COPD symptoms. Exercise in high-altitude environments can often be more uncomfortable for patients with COPD. Patients with COPD can exercise in high-altitude environments, as long as they take proper precautions.

**Unlock all 710 questions + timed mock exams**
→ https://certs.theorypractice.app/cepp
$2.99/week or $6.99/month · cancel anytime · scan to start
Unofficial study material · not affiliated with any certifying body

**7. B — Patients may need up to a year or longer to achieve full recovery from cardiac bypass surgery**
Answer: Patients may need up to a year or longer to achieve full recovery from cardiac bypass surgery. Patients typically can return to moderate exercise within a few months post-surgery, although full recovery may take up to a year or more. Care must be taken to avoid high-intensity exercises early in the rehabilitation process. Outcomes for individuals undergoing CBS vary, but gradual and progressive activity is key to recovery.

**8. B — Agility test**
Answer: Agility test Cardiovascular fitness assessment usually includes the VO2 max test, heart rate monitoring, and blood pressure measurement to evaluate the efficiency of the cardiovascular system. Agility tests, however, are part of physical fitness assessments but do not specifically measure cardiovascular fitness.

**9. C — Strength and resistance training can significantly improve muscle mass and bone density in seniors even if started late in life.**
Strength and resistance training can significantly improve muscle mass and bone density in seniors, contributing to better overall health and independence even if started later in life. It can help mitigate the effects of aging and reduce the risk of osteoporosis and sarcopenia.

**10. B — Regular aerobic exercise can significantly reduce blood pressure in hypertensive patients.**
Regular aerobic exercise has been shown to reduce both systolic and diastolic blood pressure in individuals with hypertension. This reduction is due to improvements in cardiovascular efficiency, increased nitric oxide production, and enhanced vasodilation. Strength training, when performed correctly, can also be beneficial, but it does not universally increase blood pressure. HIIT can be effective but is not the only form of beneficial exercise.

**11. B — A patient whose blood pressure remains essentially unchanged after six months of structured exercise targeted at hypertension management**
Answer: A patient whose blood pressure remains essentially unchanged after six months of structured exercise targeted at hypertension management. Those who show minimal improvement in key health parameters, despite consistent and targeted exercise programs, are often termed as nonresponders. A patient whose systolic blood pressure decreases by 15 mmHg after two months is considered a responder. A patient whose blood pressure initially drops but then rises back to pre-exercise levels after discontinuing the program for two weeks may need better adherence to the program. A patient whose overall cardiovascular fitness improves but does not enjoy the exercise sessions is responding physically but may need motivational support.

**12. C — Patients with Exercise-induced Asthma can manage their symptoms through proper warm-up routines and medication.**
Answer: Patients with Exercise-induced Asthma can manage their symptoms through proper warm-up routines and medication. It has been found that a proper warm-up routine can help in reducing the severity of asthma symptoms during exercise. Additionally, medications such as inhalers are commonly used to manage symptoms. Symptoms of Exercise-induced Asthma can be triggered by vigorous physical activity, especially in cold, dry air environments. Exercise-induced Asthma is a respiratory condition. It affects both men and women equally.

**Unlock all 710 questions + timed mock exams**
→ **https://certs.theorypractice.app/cepp**
$2.99/week or $6.99/month · cancel anytime · scan to start
Unofficial study material · not affiliated with any certifying body

**13. D — Improved cardiovascular function, increased lung capacity, enhanced overall endurance**
Answer: Improved cardiovascular function, increased lung capacity, enhanced overall endurance Aerobic exercise is known to enhance cardiovascular and respiratory health, increase endurance, and improve overall fitness levels. Reduced bone density and increased risk of joint injuries are incorrect as aerobic exercise typically promotes healthy bones and joints. Similarly, reduced muscle atrophy and worsened flexibility would not be common outcomes of regular aerobic activity. Finally, aerobic exercise generally helps in decreasing blood pressure, increasing cardiac output, and improving mental health instead of the opposite outcomes.

**14. A — Extension**
Answer: Extension The eccentric phase of a bicep curl involves elbow extension, while the concentric phase involves elbow flexion. Abduction and pronation are not motions at the elbow during a bicep curl. The eccentric phase specifically involves extension as the muscle lengthens.

**15. D — Have the client practice standing on one foot while holding onto a chair for support**
Answer: Have the client practice standing on one foot while holding onto a chair for support This exercise is effective in improving balance and stability, which can reduce the risk of falls in older adults. Running on a treadmill may be too challenging and risky for someone at high risk of falls. Lifting heavy weights can be dangerous and increase injury risk. Using loud music is not an effective method for balance training.

**16. B — Cartilage**
Answer: Cartilage Cartilage is a resilient and smooth elastic tissue; a rubber-like padding that covers and protects the ends of long bones at the joints and nerves. It is a structural component of the body that provides cushioning and support to the joints. Ligaments and tendons provide stability and attachment of muscles to bones, while muscles are responsible for movement.

**17. D — It is commonly referred to as "heel spur syndrome"**
Correct answer: It is commonly referred to as "heel spur syndrome" Plantar fasciitis typically presents as heel pain, especially with the first steps in the morning. It involves inflammation of the plantar fascia, a thick band of tissue running from the heel to the toes. "Runner's knee" commonly refers to patellofemoral pain syndrome, which affects the knee. The Achilles tendon is located at the back of the ankle, while plantar fasciitis affects the bottom of the foot. Plantar fasciitis is almost always an overuse injury, developing over weeks or months due to repetitive stress on the plantar fascia.

**18. A — An exercise where a particular muscle group is held in a fixed position, producing force without causing movement at the joint**
Answer: An exercise where a particular muscle group is held in a fixed position, producing force without causing movement at the joint Static hold exercises involve maintaining a specific posture that engages a muscle group without changing the angle of the joint. An example of this type of exercise is holding a squat position. Explosive movements pertain to power exercises. Constant speed movements relate to isokinetic exercises, which often require specialized equipment. Alternating between lifting and lowering a weight defines isotonic exercises.

**19. A — This is a normal resting heart rate.**
Answer: This is a normal resting heart rate. Resting heart rates between 60 and 100 bpm are considered normal for adults. Clinicians typically do not worry about heart rates within this range unless the patient is symptomatic. Tachycardia is defined as a resting heart rate greater than 100 bpm.

**Unlock all 710 questions + timed mock exams**
→ https://certs.theorypractice.app/cepp
$2.99/week or $6.99/month · cancel anytime · scan to start
Unofficial study material · not affiliated with any certifying body

**20. C — Sudden onset of chest pain, shortness of breath, and lightheadedness**
Answer: Sudden onset of chest pain, shortness of breath, and lightheadedness These symptoms could indicate a serious cardiovascular condition such as a heart attack or pulmonary embolism. Immediate medical attention is necessary. Mild soreness in the knees is often due to the stress of jogging and usually resolves with rest and proper care. Occasional dizziness upon standing may be due to postural changes and is usually not an emergency unless it persists. Minor lower back pain after lifting weights should be monitored but does not typically require an immediate referral.

**21. A — Hypoxia**
Answer: Hypoxia Hypoxia refers to a deficiency in the amount of oxygen reaching the tissues. During high-intensity exercise, muscle cells may not receive enough oxygen, leading to this condition. Hypercapnia is the presence of an abnormally high level of carbon dioxide in the blood. Anemia is a condition marked by a deficiency of red blood cells or hemoglobin in the blood, leading to fatigue. Ischemia is a restriction in blood supply to tissues, causing a shortage of oxygen and glucose needed for cellular metabolism.

**22. A — Ask the patient to take a deep breath, place the mouthpiece into the mouth, and blow out as hard and fast as possible into the device.**
Answer: Ask the patient to take a deep breath, place the mouthpiece into the mouth, and blow out as hard and fast as possible into the device. The purpose of the peak flow meter is to measure the maximum speed of expiration, which requires a rapid and forceful breath out. Exhaling slowly and steadily or inhaling through the device does not provide accurate measurements. The procedure needs to be standardized, capturing maximum effort for a reliable reading. Steady and slow exhalations do not measure the peak flow, which is essential for assessing lung function accurately.

**23. D — Perceived susceptibility, perceived severity, perceived benefits**
Answer: Perceived susceptibility, perceived severity, perceived benefits The Health Belief Model outlines multiple factors that influence health behaviors. It's not a single factor but a combination that determines an individual's actions. Perceived susceptibility refers to an individual's assessment of their risk of getting a condition, perceived severity is their opinion on the seriousness of the condition, and perceived benefits are the believed advantages of taking a health action.

**24. D — The patient has started asking about different types of healthy meal plans.**
Answer: The patient has started asking about different types of healthy meal plans. When a patient is starting to inquire about different ways of adopting a healthier diet, they are in the contemplation stage. In this stage, it's critical for the Clinical Exercise Physiologist (CEP) to support the patient's interest by providing information on healthy meal plans and encouraging them to take small steps towards dietary changes. Maintaining a healthy diet for over six months is characteristic of the maintenance stage. Including occasional healthy snacks is characteristic of the preparation stage. Indicating disinterest in diet change is characteristic of the precontemplation stage.

**25. C — Continuing to exercise despite injury**
Answer: Continuing to exercise despite injury Individuals with exercise addiction will often continue their exercise routines despite injuries or contraindications. This behavior can lead to further physical harm and exemplifies the compulsive nature of the addiction. Showing no interest in physical activity is generally not a sign of exercise addiction. Gaining significant weight due to inactivity might relate to other health issues but not exercise addiction. Only exercising in social settings does not encapsulate the compulsive nature of exercise addiction.

**Unlock all 710 questions + timed mock exams**
→ **https://certs.theorypractice.app/cepp**
$2.99/week or $6.99/month · cancel anytime · scan to start
Unofficial study material · not affiliated with any certifying body

**26. B — This patient may have an increased risk of falls due to decreased foot sensation.**
Answer: This patient may have an increased risk of falls due to decreased foot sensation. DPN can cause numbness in the feet of patients with diabetes. It's important to assess foot sensation and incorporate it into exercise testing and prescriptions to prevent falls and related injuries. Patients with DPN generally maintain normal upper body strength as the condition primarily affects the extremities. While diabetes can impact cognitive functions, DPN does not directly impair brain function or comprehension. Sensation recovery varies, but approximately 50% of individuals with DPN experience some degree of recovery.

**27. A — The Naughton Protocol is designed for a more gradual increase in intensity, with lower starting speeds and fewer changes in speed and grade compared to the modified Bruce Protocol.**
Answer: The Naughton Protocol is designed for a more gradual increase in intensity, with lower starting speeds and fewer changes in speed and grade compared to the modified Bruce Protocol. The Naughton Protocol starts at lower speeds and increases the intensity more gradually, making it a better choice for patients with lower exercise tolerance. In contrast, the modified Bruce Protocol increases the difficulty every minute, with both speed and grade changes. Both protocols, however, are designed to assess cardiovascular fitness and provide valuable data on a patient's exercise capacity and endurance.

**28. C — A plateau in oxygen consumption ($\dot{V}O_2$) despite an increase in workload**
Answer: A plateau in oxygen consumption ($\dot{V}O_2$) despite an increase in workload. When a plateau in $\dot{V}O_2$ occurs, it signifies that the subject has reached maximal aerobic capacity, as their body can no longer uptake oxygen more efficiently even with increased exercise intensity. Other indicators such as increasing heart rate and blood pressure do not definitively indicate that maximal aerobic capacity has been reached. A decreasing respiratory rate is contrary to the expected physiological response during increasing exercise intensity.

**29. A — Incorrect calibration of the treadmill or ergometer**
Answer: Incorrect calibration of the treadmill or ergometer Incorrect equipment calibration can lead to inaccurate measures of workload, affecting the assessment of cardiovascular fitness. For example, if the ergometer is not properly calibrated, the resistance may be higher or lower than indicated, thus altering the intensity of exercise. The graded exercise test should follow a protocol that individualizes the progression of the workload based on the subject’s fitness level to avoid premature fatigue or underestimating their fitness.

**30. D — The heart rate remains stable after reaching the anaerobic threshold.**
Answer: The heart rate remains stable after reaching the anaerobic threshold. During a maximal exercise test, as the workload increases, the heart rate also continues to increase, even past the anaerobic threshold. The anaerobic threshold is the point during intense exercise when the body can no longer meet its demand for oxygen and anaerobic metabolism is accelerated. Lactic acid levels rise, and VO2 max indicates the maximum capacity to take in, transport, and use oxygen. The RER, the ratio of carbon dioxide produced to oxygen consumed, often exceeds 1.0 at maximal exertion due to increased carbohydrate metabolism.

**Unlock all 710 questions + timed mock exams**
→ **https://certs.theorypractice.app/cepp**
$2.99/week or $6.99/month · cancel anytime · scan to start
Unofficial study material · not affiliated with any certifying body

# Ready to pass?

Unlock the full ACSM CEP Clinical Exer Prep bank, every explanation, and unlimited timed mock exams.

**Scan to start practising**
https://certs.theorypractice.app/cepp

Also on iOS & Android — search your exam name on the App Store or Google Play

**Unlock all 710 questions + timed mock exams**
→ **https://certs.theorypractice.app/cepp**
$2.99/week or $6.99/month · cancel anytime · scan to start
Unofficial study material · not affiliated with any certifying body