



NCP Nurse Competency Prep

Free Practice Test — 30 Real Exam-Style Questions

with full answer key & explanations

**Unlock the full bank of 976 questions
+ unlimited timed mock exams + mistake book**

Practice on the web: <https://certs.theorypractice.app/ncptestprep>

\$2.99 / week · \$6.99 / month · cancel anytime

What you unlock: all 976 questions • unlimited timed mock exams • mistake book • instant explanations

Also on iOS & Android — and watch the full Q&A walkthrough on [YouTube @CertsQuizPrep](#)



Unlock all 976 questions + timed mock exams

→ <https://certs.theorypractice.app/ncptestprep>

\$2.99/week or \$6.99/month · cancel anytime · scan to start



Practice Questions

Try all 30 first, then check the answer key at the back.

Want the other 946+ questions & full timed mock exams? Unlock at
<https://certs.theorypractice.app/ncptestprep>

1. A patient is ordered acetaminophen 650 mg PO. The medication is available as 325 mg tablets. How many tablets should the nurse administer?

- A. 1 tablet
- B. 1.5 tablets
- C. 3 tablets
- D. 2 tablets

2. A physician orders 1000 mL of 0.9% Normal Saline to infuse over 8 hours. What is the correct infusion rate in mL/hr?

- A. 150 mL/hr
- B. 120 mL/hr
- C. 125 mL/hr
- D. 100 mL/hr

3. An IV is ordered to infuse at 50 mL/hr using tubing with a drop factor of 15 gtt/mL. What is the correct drip rate in drops per minute?

- A. 8 gtt/min
- B. 13 gtt/min
- C. 10 gtt/min
- D. 15 gtt/min

Also on iOS & Android — and watch the full Q&A walkthrough on [YouTube](#)
[@CertsQuizPrep](#)

4. A child weighing 44 pounds is prescribed amoxicillin 20 mg/kg/day divided into three doses. What is the dose per administration in mg?

- A. 133 mg
- B. 100 mg
- C. 150 mg
- D. 120 mg



Unlock all 976 questions + timed mock exams

→ <https://certs.theorypractice.app/ncptestprep>

\$2.99/week or \$6.99/month · cancel anytime · scan to start



5. A medication order reads morphine sulfate 8 mg IM. The available vial contains 10 mg/mL. How many mL should the nurse draw up?

- A. 0.6 mL
- B. 1 mL
- C. 1.25 mL
- D. 0.8 mL

6. Convert 0.5 grams to milligrams.

- A. 5 mg
- B. 5000 mg
- C. 500 mg
- D. 50 mg

Want the other 946+ questions & full timed mock exams? Unlock at

<https://certs.theorypractice.app/ncptestprep>

7. A heparin drip is ordered at 1200 units/hr. The IV bag contains 25,000 units in 500 mL. What is the infusion rate in mL/hr?

- A. 18 mL/hr
- B. 24 mL/hr
- C. 20 mL/hr
- D. 30 mL/hr

8. A patient is to receive cefazolin 500 mg IM. The powder must be reconstituted with 2 mL of sterile water to yield 225 mg/mL. How many mL will the nurse administer?

- A. 2.2 mL
- B. 2 mL
- C. 2.5 mL
- D. 1.8 mL

9. A safe dosage range for digoxin is 0.125 to 0.25 mg daily. A physician orders 0.5 mg PO daily. What should the nurse do?

- A. Administer the medication as ordered
- B. Give half the dose
- C. Document refusal to give
- D. Question the order with the physician



Unlock all 976 questions + timed mock exams

→ <https://certs.theorypractice.app/ncptestprep>

\$2.99/week or \$6.99/month · cancel anytime · scan to start



Also on iOS & Android — and watch the full Q&A walkthrough on [YouTube](#)
[@CertsQuizPrep](#)

10. An IV of 250 mL is to infuse over 2 hours using a microdrip set (60 gtt/mL). What is the drip rate in gtt/min?

- A. 63 gtt/min
- B. 150 gtt/min
- C. 125 gtt/min
- D. 100 gtt/min

11. A patient weighs 176 pounds. The medication order is for 5 mg/kg. What is the correct total dose in mg?

- A. 880 mg
- B. 400 mg
- C. 350 mg
- D. 440 mg

12. An order reads: Give potassium chloride 30 mEq PO. Available: 20 mEq per 15 mL. How many mL should be administered?

- A. 22.5 mL
- B. 20 mL
- C. 25 mL
- D. 15 mL

Want the other 946+ questions & full timed mock exams? Unlock at
<https://certs.theorypractice.app/ncptestprep>

13. Regular insulin is ordered at 8 units/hr via infusion pump. The IV bag contains 100 units in 100 mL of 0.9% NS. What is the infusion rate in mL/hr?

- A. 10 mL/hr
- B. 6 mL/hr
- C. 12 mL/hr
- D. 8 mL/hr



Unlock all 976 questions + timed mock exams

→ <https://certs.theorypractice.app/ncptestprep>

\$2.99/week or \$6.99/month · cancel anytime · scan to start



14. Convert 2.5 liters to milliliters.

- A. 25 mL
- B. 500 mL
- C. 2500 mL
- D. 250 mL

15. A child weighing 15 kg is prescribed phenytoin 5 mg/kg/day in two divided doses. What is the amount per dose in mg?

- A. 75 mg
- B. 37.5 mg
- C. 30 mg
- D. 40 mg

Also on iOS & Android — and watch the full Q&A walkthrough on [YouTube](#)
[@CertsQuizPrep](#)

16. A medication label reads 250 mcg per tablet. The order is for 0.5 mg. How many tablets should be given?

- A. 2 tablets
- B. 1 tablet
- C. 0.5 tablet
- D. 4 tablets

17. An IV infusion of 1500 mL is ordered to run at 100 mL/hr. How many hours will it take to infuse completely?

- A. 12 hours
- B. 18 hours
- C. 10 hours
- D. 15 hours

18. A patient is to receive vancomycin 1 gram in 250 mL over 90 minutes. What is the infusion rate in mL/hr?

- A. 180 mL/hr
- B. 125 mL/hr
- C. 167 mL/hr
- D. 150 mL/hr



Unlock all 976 questions + timed mock exams

→ <https://certs.theorypractice.app/ncptestprep>

\$2.99/week or \$6.99/month · cancel anytime · scan to start



Want the other 946+ questions & full timed mock exams? Unlock at
<https://certs.theorypractice.app/ncptestprep>

19. An order reads: Administer ondansetron 0.15 mg/kg IV. The patient weighs 132 pounds. What is the correct dose in mg?

- A. 12 mg
- B. 9 mg
- C. 8 mg
- D. 10 mg

20. A patient is receiving dopamine at 12 mL/hr. The IV bag contains 400 mg in 250 mL D5W. How many mcg/min is the patient receiving?

- A. 320 mcg/min
- B. 300 mcg/min
- C. 350 mcg/min
- D. 280 mcg/min

21. Which pharmacokinetic process describes the movement of a drug from the site of administration into the bloodstream?

- A. Distribution
- B. Metabolism
- C. Excretion
- D. Absorption

Also on iOS & Android — and watch the full Q&A walkthrough on [YouTube](#)
[@CertsQuizPrep](#)

22. A drug with a half-life of 6 hours is administered at 0800. Approximately what percentage of the original dose remains in the body at 2000?

- A. 12.5%
- B. 75%
- C. 25%
- D. 50%



Unlock all 976 questions + timed mock exams

→ <https://certs.theorypractice.app/ncptestprep>

\$2.99/week or \$6.99/month · cancel anytime · scan to start



23. The therapeutic index of a drug is best defined as the relationship between which two factors?

- A. The absorption rate and the elimination rate
- B. The toxic dose and the therapeutic dose
- C. The peak level and the trough level
- D. The loading dose and the maintenance dose

24. Which term describes the percentage of an administered drug dose that reaches systemic circulation in unchanged form?

- A. Bioavailability
- B. Half-life
- C. Therapeutic index
- D. Peak concentration

Want the other 946+ questions & full timed mock exams? Unlock at
<https://certs.theorypractice.app/ncptestprep>

25. A drug that binds to a receptor but does not activate it and prevents other substances from binding is classified as which of the following?

- A. Agonist
- B. Partial agonist
- C. Prodrug
- D. Antagonist

26. The first-pass effect primarily affects drugs administered by which route?

- A. Sublingual
- B. Intramuscular
- C. Oral
- D. Intravenous

27. Which pharmacokinetic process involves the reversible transfer of a drug between body compartments such as blood, tissues, and organs?

- A. Distribution
- B. Absorption
- C. Metabolism
- D. Excretion



Unlock all 976 questions + timed mock exams

→ <https://certs.theorypractice.app/ncptestprep>

\$2.99/week or \$6.99/month · cancel anytime · scan to start



Also on iOS & Android — and watch the full Q&A walkthrough on [YouTube](#)
[@CertsQuizPrep](#)

28. A patient is receiving a drug with a narrow therapeutic index. What is the primary nursing implication for this patient?

- A. Loading doses are contraindicated
- B. The drug has minimal side effects
- C. Close monitoring is required because the margin between therapeutic and toxic doses is small
- D. The drug can be given at varying doses with minimal risk

29. A patient receiving vancomycin infusion develops flushing, pruritus, and hypotension. Which adverse reaction should the nurse suspect?

- A. Stevens-Johnson syndrome
- B. Red man syndrome
- C. Anaphylaxis
- D. Nephrotoxicity

30. A patient is prescribed warfarin therapy. Which laboratory value is most important for the nurse to monitor?

- A. Platelet count
- B. Hemoglobin
- C. White blood cell count
- D. INR



Unlock all 976 questions + timed mock exams

→ <https://certs.theorypractice.app/ncptestprep>

\$2.99/week or \$6.99/month · cancel anytime · scan to start



Answer Key & Explanations

You just practised 30 of 976. Unlock every question + timed mocks at
<https://certs.theorypractice.app/ncptestprep>

1. D — 2 tablets

Using the formula: Desired dose (650 mg) divided by Available dose (325 mg per tablet) = $650 / 325 = 2$ tablets.

2. C — 125 mL/hr

To calculate mL/hr: Total volume (1000 mL) divided by Time in hours (8 hours) = $1000 / 8 = 125$ mL/hr.

3. B — 13 gtt/min

Formula: (mL/hr × drop factor) / 60 minutes = $(50 \times 15) / 60 = 750 / 60 = 12.5$, rounded to 13 gtt/min.

4. A — 133 mg

First convert pounds to kg: $44 \text{ lb} / 2.2 = 20 \text{ kg}$. Daily dose: $20 \text{ kg} \times 20 \text{ mg/kg} = 400 \text{ mg/day}$. Per dose: $400 \text{ mg} / 3 \text{ doses} = 133.3 \text{ mg}$, rounded to 133 mg per dose.

5. D — 0.8 mL

Using the formula: Desired dose (8 mg) divided by Available concentration (10 mg/mL) = $8 / 10 = 0.8$ mL.

6. C — 500 mg

To convert grams to milligrams, multiply by 1000. Therefore, $0.5 \text{ g} \times 1000 = 500 \text{ mg}$.

7. B — 24 mL/hr

Using ratio-proportion: $25,000 \text{ units} : 500 \text{ mL} = 1200 \text{ units} : X \text{ mL}$. Cross multiply: $25,000X = 600,000$. $X = 600,000 / 25,000 = 24$ mL/hr.

8. A — 2.2 mL

After reconstitution, the concentration is 225 mg/mL. Using the formula: Desired dose (500 mg) divided by Available concentration (225 mg/mL) = $500 / 225 = 2.22$ mL, rounded to 2.2 mL.

9. D — Question the order with the physician

The ordered dose of 0.5 mg exceeds the maximum safe dose of 0.25 mg daily (it is double the maximum). The nurse should question this order with the physician before administering to prevent toxicity.

10. C — 125 gtt/min

For microdrip (60 gtt/mL), the drip rate equals the hourly rate. First find mL/hr: $250 \text{ mL} / 2 \text{ hr} = 125 \text{ mL/hr}$. With microdrip, $125 \text{ mL/hr} = 125 \text{ gtt/min}$.

11. B — 400 mg

First convert pounds to kg: $176 \text{ lb} / 2.2 = 80 \text{ kg}$. Then calculate dose: $80 \text{ kg} \times 5 \text{ mg/kg} = 400 \text{ mg}$.

12. A — 22.5 mL

Using ratio-proportion: $20 \text{ mEq} : 15 \text{ mL} = 30 \text{ mEq} : X \text{ mL}$. Cross multiply: $20X = 450$. $X = 450 / 20 = 22.5$ mL.

13. D — 8 mL/hr

The concentration is 100 units in 100 mL = 1 unit/mL. To deliver 8 units/hr: $8 \text{ units} / 1 \text{ unit per mL} = 8 \text{ mL/hr}$.



Unlock all 976 questions + timed mock exams

→ <https://certs.theorypractice.app/ncptestprep>

\$2.99/week or \$6.99/month · cancel anytime · scan to start

Unofficial study material · not affiliated with any certifying body



14. C — 2500 mL

To convert liters to milliliters, multiply by 1000. Therefore, $2.5 \text{ L} \times 1000 = 2500 \text{ mL}$.

15. B — 37.5 mg

Calculate total daily dose: $15 \text{ kg} \times 5 \text{ mg/kg} = 75 \text{ mg/day}$. Divide by 2 doses: $75 \text{ mg} / 2 = 37.5 \text{ mg per dose}$.

16. A — 2 tablets

First convert 0.5 mg to mcg: $0.5 \text{ mg} \times 1000 = 500 \text{ mcg}$. Then: $500 \text{ mcg} / 250 \text{ mcg per tablet} = 2 \text{ tablets}$.

17. D — 15 hours

Total volume (1500 mL) divided by infusion rate (100 mL/hr) = $1500 / 100 = 15 \text{ hours}$.

18. C — 167 mL/hr

Convert 90 minutes to hours: $90 / 60 = 1.5 \text{ hours}$. Then calculate rate: $250 \text{ mL} / 1.5 \text{ hr} = 166.7 \text{ mL/hr}$, rounded to 167 mL/hr.

19. B — 9 mg

Convert weight to kg: $132 \text{ lb} / 2.2 = 60 \text{ kg}$. Calculate dose: $60 \text{ kg} \times 0.15 \text{ mg/kg} = 9 \text{ mg}$.

20. A — 320 mcg/min

First find mg/hr: $(400 \text{ mg} / 250 \text{ mL}) \times 12 \text{ mL/hr} = 19.2 \text{ mg/hr}$. Convert to mcg/min: $(19.2 \text{ mg} \times 1000 \text{ mcg/mg}) / 60 \text{ min} = 19,200 / 60 = 320 \text{ mcg/min}$.

21. D — Absorption

Absorption is the process by which a drug moves from its site of administration (such as the GI tract, muscle, or skin) into the systemic circulation. This is the first phase of pharmacokinetics.

22. C — 25%

After one half-life (6 hours, at 1400), 50% remains. After two half-lives (12 hours, at 2000), 25% of the original dose remains. Each half-life reduces the drug concentration by half.

23. B — The toxic dose and the therapeutic dose

The therapeutic index is the ratio between the dose that produces toxicity and the dose that produces a therapeutic effect. A wider therapeutic index indicates a safer drug with a larger margin between effective and toxic doses.

24. A — Bioavailability

Bioavailability refers to the fraction of an administered dose that reaches the systemic circulation unchanged and is available to produce its therapeutic effect. Intravenous drugs have 100% bioavailability, while oral drugs may have lower bioavailability due to first-pass metabolism.

25. D — Antagonist

An antagonist binds to a receptor without activating it and blocks the binding of agonists or endogenous ligands, thereby preventing receptor activation and the associated biological response.

26. C — Oral

The first-pass effect occurs when oral drugs are absorbed from the GI tract into the portal circulation and metabolized by the liver before reaching systemic circulation. This can significantly reduce the bioavailability of orally administered drugs.



Unlock all 976 questions + timed mock exams

→ <https://certs.theorypractice.app/ncptestprep>

\$2.99/week or \$6.99/month · cancel anytime · scan to start

Unofficial study material · not affiliated with any certifying body



27. A — Distribution

Distribution is the process by which a drug is transported from the bloodstream to various tissues and organs throughout the body. Factors affecting distribution include blood flow, protein binding, and tissue permeability.

28. C — Close monitoring is required because the margin between therapeutic and toxic doses is small

A narrow therapeutic index means there is a small margin between therapeutic and toxic doses. Close monitoring of drug levels and patient response is essential to prevent toxicity while maintaining therapeutic effectiveness.

29. B — Red man syndrome

Red man syndrome is a vancomycin-related adverse effect characterized by flushing, pruritus, hypotension, and sometimes chest pain. It occurs due to histamine release when the drug is infused too rapidly.

30. D — INR

INR (International Normalized Ratio) is the laboratory value used to monitor warfarin effectiveness and adjust dosing. Therapeutic range is typically 2-3 for most indications.



Unlock all 976 questions + timed mock exams

→ <https://certs.theorypractice.app/ncptestprep>

\$2.99/week or \$6.99/month · cancel anytime · scan to start



Ready to pass?

Unlock the full NCP Nurse Competency Prep bank, every explanation, and unlimited timed mock exams.

Scan to start practising

<https://certs.theorypractice.app/ncptestprep>

Watch the full video walkthrough on YouTube @CertsQuizPrep



Unlock all 976 questions + timed mock exams

→ <https://certs.theorypractice.app/ncptestprep>

\$2.99/week or \$6.99/month · cancel anytime · scan to start